

Other Medical Geology Issues

Robert B. Finkelman
U.S. Geological Survey
rbf@usgs.gov



Medical Geology-Range of Issue

- Trace Element Exposure- As, Hg, F, Se, Zn, Al
- Dust- Asbestos, African, Valley Fever, Silicosis, CWP,VOG
- Organics - VOCs, MTBE, PAHs, Antibiotics, Pesticides
- Radionuclides - Radon, Radium, Uranium
- Microbes, Pathogens - West Nile Encephalitis, LaCrosse Encephalitis, Plague, Hantavirus, Rift Valley Fever, Lyme disease, etc.
- Global Climate Change

Other Medical Geology Issues – Outline

- Volcanism
- Organics (BEN)
- Radioactivity
- Pathogens and Microbes
- Occupational Health

The health effects of tephra dispersal




Volcanic tephra dispersal

- Mucous membrane irritation
- Silicosis
- Adsorbed toxins

- Calcium fluorosilicate (CaSiF_6)



**Silicotic nodule in
the lung tissue
with disruption of
surrounding
alveoli**







The health effects of volcanic gas emissions



Inert asphyxiants

- Carbon dioxide, CO₂



Irritant gases

- **Hydrofluoric acid, HF /hydrochloric acid, HCl**
 - Mucosal irritation
 - Cutaneous burns
 - Respiratory disease
- **Sulphur dioxide, SO₂**
 - Asthma
 - Acid rain



Noxious asphyxiants

- **Hydrogen sulphide, H₂S**

7 µg/m³ – ‘rotten egg’ smell

15,000 µg/m³ – eye irritation

480,000 µg/m³ – risk of pulmonary oedema

1,500,000 µg/m³ – lethal

Health effects of other eruptive events

- **Lava flows**
- **Pyroclastic flows**
- **Volcanic activity and aquatic environments**




Medical – geology interface

- **A review of recent disaster responses has indicated a degree of mismatching between the acquisition of eruption data and its health-related utilisation.**
- **From a medical perspective, geologic monitoring has a pivotal - but often under-utilised – role in helping those living in the shadows of volcanoes.**




Pesticides almost always occur as mixtures



BALKAN ENDEMIC NEPHROPATHY (BEN)










Radionuclide Samples Collected from Public Water Supplies


Percent of samples exceeding targeted concentrations




PREDICTED CANCER FATALITIES DUE
TO IONIZING RADIATION:
GENERAL POPULATION: AVERAGE DOSE


	mrem/yr	RADIATION FATALITIES	
		Total Number in U.S. per year	Per Million Persons per year
MEDICAL DIAGNOSTIC	70	3080	14
COSMIC RADIATION	35	1540	7
TERRESTRIAL (rocks and soil, etc.)	35	1540	7
POTASSIUM-40 IN FOOD	20	880	4
NUCLEAR WEAPONS FALLOUT	4.4	194	0.9
USE OF NATURAL GAS IN HOMES	2	89	0.4
BURNING OF COAL	1	44	0.2
SLEEPING WITH ANOTHER PERSON	0.1	4.4	0.02
NUCLEAR POWER	0.1	4.4	0.02
CONSUMER PRODUCTS (TV, etc.)	0.03	1.3	0.006
TOTAL	168	7377	

Typical Range of Uranium concentration in coal, fly ash, and a variety of common rocks





Photograph of hollow
glassy fly ash particle
(0.01 cm D)



Fission track
radiograph of the same
particle

Geographic Analysis of Disease Risk

- Where are the potential areas of disease?
- Who are the populations at risk now and in the future?
- When might an outbreak occur?
- How can outbreaks be mitigated?



Landscape Epidemiology

By knowing the ecological conditions necessary for the maintenance of specific pathogens in nature, one can use these characteristics to identify the spatial and temporal distribution of disease risk.




Locating Mosquito Breeding Sites

Use land characteristics, FEMA flood maps and imagery to identify locations of potential *Culiseta melanura* habitat, but still accessible by roads or trails, where mosquito traps may be placed; determine risk to human health.



Valley Fever: Geological/Ecological occurrence modeling



← geology

← geochemistry

← soils

← remote sensing
interpretation

← elevation

....


← spatial data_i



← *Valley Fever favorableness*

Analysis of Lyme Disease


LYME DISEASE — reported cases*, United States, 1997



*One dot=one case, randomly placed within county of residence.

In 1997, a total of 12,801 cases of Lyme disease were reported by 45 states and the District of Columbia. The 10 states with the highest incidence of Lyme disease cases per 100,000 population were Connecticut, Rhode Island, New Jersey, New York, Pennsylvania, Delaware, Massachusetts, Wisconsin, Minnesota, and Maryland. These states accounted for 52% of the reported Lyme disease cases in 1997.

Established* and reported** distribution of the Lyme disease vectors *Ixodes scapularis* (*I. dammini*) and *Ixodes pacificus*, by county, United States, 1907-1996




Vector presence
 Established *Ixodes scapularis*
 Reported *Ixodes scapularis*
 Established *Ixodes pacificus*
 Reported *Ixodes pacificus*

CDC
 Centers for Disease Control and Prevention

at least 6 ticks or 2 life stages (larvae, nymphs, adults) identified.
 at least 1 tick identified


National Lyme disease risk map with four categories of risk





Areas of predicted Lyme disease transmission
 High risk
 Moderate risk
 Low risk
 Minimal or no risk

Note: This map demonstrates an approximate distribution of predicted Lyme disease risk in the United States. The true relative risk in any given county compared with other counties might differ from that shown here and might change from year to year. Risk categories are defined in the accompanying text. Information on risk distribution within states and counties is best obtained from state and local public health authorities.

Spatial Forecast of Lyme Disease



Plague Cases in the SW United States



#65 - Mixed Rangeland


#82 - Desert Shrubland & Grassland

#148 - Ponderosa Pine

#184 - Pinyon Juniper

Spread of West Nile Virus 1999-2002

(Avian Mortality)



Conclusions



- Geographic analysis tools can model the processes that affect the occurrence and spread of diseases
- Collaborating with health professionals, we can understand the linkages between environmental factors and human health and work to reduce the risk of disease.

Medical Geology and Occupational Health

- Hard Rock Mining
- Coal Mining
- Asbestos Mining and Processing
- Ore Processing
- Farming
- Power Plant Workers

MINING AND OCCUPATIONAL HEALTH



Immediate and short term health effects

- **Trauma** eg cave-ins and other accidents, including explosions
- **Thermal injury**
- **Pressure effects**
- **Toxic gas inhalation**
- **Injury to sensory organs** (noise –induced hearing loss; ear, nose and throat and visual irritation)

Delayed /chronic health effects

- **Carcinogenicity**
- **Dermatological effects**
- **Respiratory effects**



Carcinogenicity

EXAMPLES OF PROBABLE OR DEFINITE CARCINOGENS ASSOCIATED WITH MINING / SMELTING

Asbestos

Coke oven emissions


Uranium and radon

Benzene


Nickel

Arsenic

Lung diseases associated with mining 1: exposure to coal dust




Complicated coal workers pneumoconiosis





Progressive massive pulmonary fibrosis in a coal worker

Lung diseases associated with mining 2: the spectrum of silica-related disorders



Increased risk of protracted TB






Crocidolite (Blue Asbestos)


Lung diseases associated with mining 3: the spectrum of asbestos-related disorders




Asbestos fibre in lung tissue



Asbestosis



Enormous mesothelioma tumour mass filling chest cavity



Lung cancer adjacent to main bronchus

Mercury in the soil and food chain



Mercury poisoning causes motor and visual impairment

Mseleni Joint Disease

- Multiple epiphyseal displasia (long bones have malformed growth)
- Polyarticular osteoarthritis (arthritis of several joints)
- Protrusio acetabuli (hip disorder)
- Dwarfism

Disease Progression



Prevalence



- Onset unknown
- Overall 39% women, 11% men
- >19, 66% women, 25% men

Prior Geochemical Research

- Soils

- Deficient: N, P, K, S, Ca, Zn, Cu, and B
- Suspected: Mo
- Not studied: F, I, V and Se

Grey Fernwood Sand

- Near neutral $\text{pH}_{\text{H}_2\text{O}}$ 6.9
- < 4% clay (kaolinite and quartz)
- Low organic C ~1.6%
- CEC 2.0 $\text{cmol}_c \text{ kg}^{-1}$



